## INTERVENTIONAL SCHEDULING



Radiology Ltd. Interventional Imaging Services

Interventional Scheduling Tel: (520) 545-1906 Fax: (520) 545-1898

### The state of the insurance of the in
TIENT IS ON THE FOLLOWING ANTICOAGULANT(S)  Aggrenox for 5 days  Aspirin for 5 days  Brilinta for 5 days  Brilinta for 5 days  Coumadin for 5 days  Effent for 7 days  Effent for 7 days  Eliquis for 48 hours  Cother:  Patient may discontinue above indicated  Anticoagulant(s) 5-7 days before procedure.
Middle Initial)
Middle Initial)   BEST TIME TO CONTACT: AM PM
BEST TIME TO CONTACT: AM PM  INS. PHONE: ()  ould you like authorization assistance? □ YES □ NO  ation and a copy of the insurance card with the order.  CareMore, Humana and United Healthcare insurance plans.  ATIENT IS ON THE FOLLOWING ANTICOAGULANT(S) □ Aggrenox for 5 days □ Plavix for 5 days □ Aspirin for 5 days □ Pletal □ Brilinta for 5 days □ Pradaxa for 2 days with normal renal function □ Coumadin for 5 days □ Pradaxa for 5 days with normal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 2 days with abnormal renal function □ Pradaxa for 5 days □ Coumadin for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 2 days with abnormal renal function □ Pradaxa for 5 days □ Coumadin for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 2 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with abnormal renal function □ Pradaxa for 5 days with
INS. PHONE: (
and a copy of the insurance card with the order.  CareMore, Humana and United Healthcare insurance plans.  ITIENT IS ON THE FOLLOWING ANTICOAGULANT(S)  Aggrenox for 5 days Aspirin for 5 days Brilinta for 5 days Coumadin for 5 days Effient for 7 days Eliquis for 48 hours Lovenox for 24 hours Other: Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
Aggrenox for 5 days Aspirin for 5 days Brilinta for 5 days Coumadin for 5 days Effient for 7 days Eliquis for 48 hours Lovenox for 24 hours Other: Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
Aggrenox for 5 days Aspirin for 5 days Brilinta for 5 days Coumadin for 5 days Effient for 7 days Eliquis for 48 hours Lovenox for 24 hours Other: Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
TIENT IS ON THE FOLLOWING ANTICOAGULANT(S)  Aggrenox for 5 days Plavix for 5 days Pletal Pradaxa for 2 days with normal renal function Pradaxa for 5 days With normal renal function Pradaxa for 5 days with normal renal function Warfarin for 5 days Warfarin for 5 days Lovenox for 24 hours  Other:  Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
Aggrenox for 5 days Aspirin for 5 days Brilinta for 5 days Coumadin for 5 days Effient for 7 days Eliquis for 48 hours Lovenox for 24 hours Other: Patient may discontinue above indicated Anticoagulant(s) 5-7 days Pletal Pradaxa for 2 days with normal renal function Pradaxa for 5 days with normal renal function Warfarin for 5 days Warfarin for 5 days Xarelto for 24 hours
□ Brilinta for 5 days □ Coumadin for 5 days □ Effient for 7 days □ Eliquis for 48 hours □ Lovenox for 24 hours □ Other: □ Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure. □ Pradaxa for 2 days with normal renal function □ Pradaxa for 5 days with normal renal function □ Warfarin for 5 days □ Xarelto for 24 hours
Coumadin for 5 days  Effient for 7 days  Eliquis for 48 hours  Lovenox for 24 hours  Other:  Patient may discontinue above indicated  Anticoagulant(s) 5-7 days before procedure.
☐ Eliquis for 48 hours ☐ Warfarin for 5 days ☐ Lovenox for 24 hours ☐ Xarelto for 24 hours ☐ Other: ☐  Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
□ Lovenox for 24 hours □ Xarelto for 24 hours □ Other: □ Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
Patient may discontinue above indicated Anticoagulant(s) 5-7 days before procedure.
Anticoagulant(s) 5-7 days before procedure.
Tes Tivo Di. Illidas
*Hospital-based exams are performed by
Radiology Ltd. physicians in a hospital setting. If exam is deemed hospital-based, it will
be forwarded to one of the hospitals where
Radiology Ltd. physicians perform the exam.
HOSPITAL SCHEDULING
er Procedures)  St. Joseph's Hospital
Scheduling - Tel: 872-7200 Fx: 872-7884
Cath Lab/IR - Tel: 873-3801 Fx: 751-5139
Tucson Medical Center
Scheduling - Tel: 324-2075 Fx: 324-6162
Cath Lab - Tel: 324-5034 Fx: 324-5022
Where:
Date:
uled Appointment:  ME: AM PM

The information contained in this facsimile message is CONFIDENTIAL and/or LEGALLY PRIVILEGED information intended only for the use of the facility named above. If you have received this in error, please call (520) 545-1969.

If you have any questions, please call: (520) 545-1906